DOCUMENT RESUME

ED 467 260 CG 031 879

AUTHOR Magyar-Moe, J. L.; McDermott, D.; Edwards, L. M.; Pedrotti,

J. T.; LaRue, S.; Stone, M. E.; Spalitto, S. V.; Diamond, K.

L.

TITLE Variables Related to Life Satisfaction among Senior Adults in

Assisted Living Facilities.

PUB DATE 2001-09-16

NOTE 15p.

PUB TYPE Reports - Research (143)

EDRS PRICE EDRS Price MF01/PC01 Plus Postage.

DESCRIPTORS Depression (Psychology); *Life Satisfaction; Mental Health;

*Older Adults; *Personal Care Homes; *Predictor Variables;

Spirituality

IDENTIFIERS Hope

ABSTRACT

As the population of the United States rapidly grays, the plight of older people living in assisted living situations is of growing concern to health and mental health care teams, as well as the families of the individuals residing in these living environments. This large group of residents, past the stage of living in their own homes, yet not in need of skilled care, is at risk for low satisfaction with life, depression, and hopelessness. Due to the potentially positive roles of hope and spirituality in lowering depression levels and raising life satisfaction ratings, the current study was conducted in order to empirically examine the relationship among these variables for senior adults in assisted living facilities. Results reveal that life satisfaction correlated positively with hope and spirituality, and negatively with depression. Life satisfaction did not correlate significantly with indicators of poor health as expected, however, the correlation did approach significance. Additionally, depression was found to be significantly negatively correlated with spirituality and hope. (Contains 27 references.) (Author/GCP)



Running head: VARIABLES RELATED TO LIFE SATISFACTION

Variables Related to Life Satisfaction among Senior Adults in Assisted Living Facilities

Magyar-Moe, J. L., McDermott, D., Edwards, L. M., Pedrotti, J. T., LaRue, S., Stone, M. E., Spalitto, S. V., and Diamond, K. L.

University of Kansas

U.S. DEPARTMENT OF EDUCATION Office of Educational Research and Improvement EDUCATIONAL RESOURCES INFORMATION

CENTER (ERIC)

This document has been reproduced as received from the person or organization originating it.

☐ Minor changes have been made to improve reproduction quality.

Points of view or opinions stated in this document do not necessarily represent official OERI position or policy.

PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL HAS BEEN GRANTED BY



Abstract

As the population of the United States rapidly grays, the plight of older people residing in assisted living situations is of growing concern to health and mental health care teams, as well as the families of the individuals residing in these living environments. This large group of residents, past the stage of living in their own homes, yet not in need of skilled care, is at risk for low satisfaction with life, depression, and hopelessness. Due to the potentially positive roles of hope and spirituality in lowering depression levels and raising life satisfaction ratings, the current study was conducted in order to empirically examine the relationship among these variables for senior adults in assisted living facilities.



Variables Related to Life Satisfaction among Senior Adults in Assisted Living Facilities

As the population of the United States rapidly grays, the plight of older people residing in assisted living situations is of growing concern to health and mental health care teams, as well as the families of the individuals residing in these living environments. This large group of residents, past the stage of living in their own homes, yet not in need of skilled care, is at risk for low satisfaction with life, depression, and hopelessness. Senior adults who live in their own homes are relatively free to structure their lives as they wish, while patients in nursing homes are tended by a team of health professionals, and thus have many of their social as well as physical needs met. Those elderly in assisted living facilities, however, seem to exist in a limbo past retirement, but not in need of skilled nursing care. Despite the less than optimal conditions experienced in many assisted living situations, some people maintain, even thrive, while others decline. The purpose of the present research was to explore variables that are related to life satisfaction among senior adult residents in assisted living communities. Using life satisfaction as the criterion variable, we hypothesized that results would show a negative relationship with depression and symptoms of poor health, and a positive relationship with hope and spirituality. **Depression Rates in Senior Populations**

The prevalence of mental disorders found among elderly populations appears to be similar to that found in younger populations with the exception of depression and dementia which are found more often in senior adults (Rogers, 1999). According to McCullough (1991) and Molinari (1991), depression is found in approximately .7% to 2.7% of the general population



whereas depressive symptoms among the elderly are estimated to range from 2.5% to 45% of the population with an average of 20%. More specifically, Oxman, Barrett, Barrett, and Gerber (1990) report 6% to 9% prevalence estimates for major depressive disorder diagnoses of elderly adults treated in primary care facilities, and Parmelee, Katz, and Lawton (1992) report 10 % - 15% prevalence rates of major depression in elders with chronic medical conditions and in senior adults who reside in institutional settings. Older adults who do not meet full criteria for major depression may receive a diagnosis of minor depression which has prevalence estimates ranging from 4% to 26% for elders living in the community (Blazer, Hughes, & George, 1987), 47% to 52% for elderly medical outpatients (Oxman et al.), and 18% to 30.5% of senior residents in institutional settings (Parmelee et al.).

Depression and Health Status

Maintaining optimal physical health is a salient issue among many senior adults. Rogers (1999) reports that mental illness, specifically depression, can significantly interfere with older adult's abilities to cope with and recover from physical conditions. In addition, lower rates of recovery from depression have been linked with comorbid medical problems and long-term disability (Rogers). Gerety and Farnett (1995) report that physical illness and depression may co-occur in up to 45% of individuals with physical ailments in senior populations.

A number of self-report survey studies with elderly individuals have concluded that among this population, depressive symptoms are significantly related to medical problems and problems with daily living skills. Those who perceived their health as poor and those who did not feel they had control over their health also had more depressive symptoms. Self-reported



ailing health, inability to participate fully in daily living activities, and symptoms of physical conditions such as pain and arthritis were all associated with higher levels of depression among those surveyed (Henderson et al., 1993; Kennedy et al., 1989).

The Role of Hope

C. R. Snyder (1994) defines hope as an active cognitive process of goal selection, empowerment, and the ability to find alternate ways to reach goals when problems are encountered. Cheavens and Gum (2000) contend that hope plays a significant role in the lives of all people, including those over age 65. They report that life satisfaction and successful aging are linked with high levels of hope in the later years whereas elderly individuals with lower levels of hope are likely to be less satisfied with life and to experience increased disease states.

Snyder and colleagues (1991) have also found strong negative relationships between hope and depression such that those individuals who are low in hope are also more depressed than those with high hope levels. Furthermore, Snyder, LaPointe, Crowson, and Early (in press) report that the thinking patterns of depressed persons and those low in hope are similar. Based on this evidence, Klausner, Snyder, and Cheavens (2001) developed a hope-based, goal focused group psychotherapy intervention for the purpose of treating individuals with late-life depression. The intervention was tested and shown to reduce depressed mood, anxiety, and hopelessness while increasing hope and social functioning. The authors concluded that these improvements have the potential to enhance the quality of life and life satisfaction in the senior population.



The Role of Spirituality

Spirituality has been postulated as one of the factors that may help some older adults to cope and even thrive despite less than ideal health status, economic status, living conditions, role loss, and loss of loved ones (Neill & Kahn, 1999). Hunsberger (1985) discovered a positive relationship between life satisfaction and religion among senior adults, and Morse and Wisocki (1988) reported levels of religious activity and beliefs to be associated with increased psychological health in the elderly.

Based on the above literature related to high levels of depression and low levels of life satisfaction among the senior population, there appears to be a need for mental health practitioners to provide interventions to senior adults to help decrease depression and increase satisfaction among these clients. Due to the potentially positive roles of hope and spirituality in lowering depression levels and raising life satisfaction ratings, the current study was conducted in order to empirically examine the relationship among these variables for senior adults in assisted living facilities.

Method

Participants

The participants in this study were 29 women and 13 men between the ages of 55 and 98 residing in Christian-affiliated assisted living facilities in two midwestern communities. The majority of participants were Caucasian and of middle to lower middle socioeconomic status.



Materials

The Geriatric Depression Scale (GDS, Yesavage et al., 1983) is a 30-item self-report measure, yielding possible scores of 0 to 30. Internal reliability studies have reported alpha levels of .88 to .94. Test-retest correlations, at intervals from 2 hours to 1 month, were found to be high, ranging from .85 to .94. Studies of construct validity have demonstrated that the GDS correlates positively with other measures of depression. The GDS has not been found to correlate with cognitive screening tests, supporting its divergent validity (Stiles & McGarrahan, 1998).

In order to measure health problems, the Symptom Checklist (SC), a 20-item instrument designed to measure frequency of psychiatric/medical symptoms was used. The SC has excellent internal consistency with alphas that range from .90 to .93, and good predictive validity has been reported (Bartone, Ursano, Wright, & Ingraham, 1989).

The Adult Hope Scale (Snyder et al, 1991) is a 12-item self report assessment designed to measure the extent to which people believe they can find ways to solve problems, and the amount of mental energy they have to pursue their goals. The scale yields a total hope score, and scores on the two sub-scales of pathways and agency. Cronbach alphas for the total hope score are reported to range from .74 to .84 and test-retest reliability over a 10-week interval was found to be high with a correlation of .80. Construct validity has been found with positive correlations between the Hope Scale and measures of optimism, expectancy for attaining goals, amount of expected control, and self-esteem. Likewise, negative correlations have been reported between the Hope Scale and measures of depression and hopelessness.



The Spiritual Well-being Scale (Ellison & Paloutzian, 1982) is a 20-item self-report instrument developed as a general indicator of the subjective states of religious and existential well-being and overall life satisfaction. The test-retest reliability coefficient is .93 and internal consistency is .89. The validity is reported as being acceptable with positive correlations between the SWB scale and standard indicators of well-being such as positive self-concept, finding meaning and purpose in life, high assertiveness, low aggressiveness, good physical health, and good emotional adjustment. The SWB scale negatively correlates with indicators of ill health, loneliness, emotional maladjustment, and dissatisfaction with life.

The Life Satisfaction Index-Z (Wood, Wylie, & Sheafor, 1969), a 13-item scale with scores ranging from 0-26, was also administered. Higher scores indicate higher life satisfaction. The split-half reliability coefficient is .79 and the coefficient alpha is .80. The Life Satisfaction Index-Z has been shown to demonstrate discriminant validity between individuals with low and high satisfaction (Neugarten, Havighurst, & Tobin, 1961).

Procedure

All data were collected in the participants' assisted living communities. Approximately two-thirds of the participants were interviewed individually by a member of the research team, while the remaining third completed the measures on their own with a researcher nearby to answer questions.

Results

To examine the extent to which depression, symptoms of poor health, hope, and spirituality were related to life satisfaction, Pearson product moment correlation coefficients



were calculated. Correlations revealed a significant negative relationship between life satisfaction and depression (r = -.748, p < .01) and a non-significant negative relationship between life satisfaction and symptoms of poor health (r = -.299). Positive, significant relationships were found for life satisfaction and hope (r = .575, p < .01), life satisfaction and each of the hope subscales (agency: r = .620, p < .01; pathways: r = .430, p < .01) and life satisfaction and spirituality (r = .497, p < .01). See Table 1 for the complete correlation matrix results.

Discussion

Life satisfaction correlated positively with hope and spirituality, and negatively with depression. Life satisfaction did not correlate significantly with indicators of poor health as expected, however, the correlation did approach significance at the .05 level. Those individuals with higher hope and spirituality showed higher life satisfaction. Additionally, depression was found to be significantly negatively correlated with spirituality, hope, and both of the hope subscales. Positive correlations were found between spirituality and hope, including both hope subscales.

Hope was a factor of major interest in the study. Strategies for teaching the hope process have been developed and used successfully with children and adults (McDermott & Snyder, 1999; Snyder 2000). Once the factors that predict life satisfaction among the assisted living population have been identified, hope building strategies for these senior adults can be developed and implemented. This research is a preliminary step toward the development of a program to enhance life satisfaction among the assisted living population.



References

Bartone, P. T., Ursano, R. J., Wright, K. M., & Ingraham, L. H. (1989). The impact of a military air disaster on the health of assistance workers. <u>Journal of Nervous and Mental Disease</u>, 177, 317-328.

Blazer, D. G., Hughes, D. C., & George, L. K. (1987). The epidemiology of depression in an elderly community population. The Gerontologist, 27, 281-287.

Brink, T. L., Yesavage, J., Heersema, P., Lum. O., Adey, M., Rose, T. L., Leirer, V. O., & Huang, V. (no date). Geriatric Depression Scale. Redlands, CA: T. L. Brink, Ph.D.\

Cheavens, J., & Gumm, A. (2000). Gray power: Hope for the ages. In C. R. Snyder (Ed.), Handbook of hope: Theory, measures, and applications, (pp. 201-221). San Diego: Academic Press.

Duke University Department of Community and Family Medicine (1979). <u>Duke-UNC</u>

<u>Health Profile.</u> Duke University Medical Center, Durham, NC.

Ellison, C. W. & Paloutzian, R. F. (1982). Spiritual Well-being Scale. Nyack, NY: Life Advance.

Gerety, M. B., & Farnett, L. (1995). Management of depression in the elderly. <u>Strategies</u> in <u>Geriatrics</u>, 2, 137-143.

Henderson, A. S., Jorm, A. F., Mackinnon, A., Christensen, H., Scott, L. R., Korten, A. E., & Doyle, C. (1993). The prevalence of depressive disorders and the distribution of depressive symptoms in later life: A survey using Draft ICD-10 and DSM-III-R. <u>Psychological Medicine</u>, 23, 719-729.



Hunsberger, B. (1985). Religion, age, life satisfaction, and perceived sources of religiousness: A study of older persons. <u>Journal of Gerontology</u>, 40, 615-620.

Kennedy, G. J., Kelman, H. R., Thomas, C., Wisniewski, W., Metz, H., & Bijor, P. E. (1989). Hierarchy of characteristics associated with depressive symptoms in an urban elderly sample. American Journal of Psychiatry, 146, 220-225.

Klausner, E. J., Snyder, C. R., & Cheavens, J. (2001). A hope-based treatment for depressed older adult patients. In G. Williamson, J. Parmelee, & D. Shaffer (Eds.), <u>Handbook on aging: Theory, research, and applications</u>. New York: Plenum Press.

McCullough, P. K. (1991). Geriatric depression: Atypical presentations, hidden meanings. Geriatrics, 46, 72-76.

McDermott, D. & Snyder, C. R. (1999). <u>Making hope happen: A workbook for turning possibilities into realities.</u> Oakland, CA: New Harbinger.

Molinari, V. A. (1991). Mental health issues in the eldery. The mentally impaired elderly: Strategies and interventions to maintain function (Special issue). Physical and Occupational Therapy in Geriatrics, 9, 23-30.

Morse, C. K., & Wisocki, P. A. (1988). Importance of religiosity to elderly adjustment.

<u>Journal of Religion and Aging, 4, 15-26.</u>

Neill, C. M., & Kahn, A. S. (1999). The role of personal spirituality and religious social activity on the life satisfaction of older widowed women. <u>Sex Roles</u>, 40, 319-329.

Neugarten, B. L., Havighurst, R. J., & Tobin, S. S. (1961). The measurement of life satisfaction. <u>Journal of Gerontology</u>, 16, 134-143.



Oxman, T. E., Barrett, J. E., Barrett, J., & Gerber, P. (1990). Symptomology of late-life minor depression among primary care patients. <u>Psychosomatics</u>, 31, 174-180.

Parmelee, P. A., Katz, I. R., & Lawton, M. P. (1992). Incidence of depression in long-term care settings. <u>Journal of Gerontology: Medical Sciences</u>, 47, M189-M196.

Rogers, A. (1999). Factors associated with depression and low life satisfaction in the low-income, frail elderly. <u>Journal of Gerontological Social Work</u>, 31, 167-194.

Snyder, C. R. (2000). <u>Handbook of hope: Theory, measures, and applications.</u> San Diego: Academic Press.

Snyder, C. R. (1994). The psychology of hope: you can get there from here. New York, NY: Free Press.

Snyder, C. R. Harris, C., Anderson, J. R., Holleran, S. A., Irving, L. M., Sigmon, S. T., Yoshinobu, L. R., Gibb, J., Langelle, C., Harney, P. (1991). The will and the ways: Development of an individual-differences measure of hope. <u>Journal of Personality and Social Psychology</u>, 60, 570-585.

Snyder, C. R., LaPointe, A. B., Crowson Jr., J. J., & Early, S. (in press). Preferences of high- and low-hope people for self-referential feedback. <u>Cognition and Emotion</u>.

Stiles, P. G., & McGarrahan, J. F. (1998). The Geriatric Depression Scale: A comprehensive review. <u>Journal of Clinical Geropsychology</u>, 4, 89-110.

Wood, V., Wylie, M. F., & Sheafor, B. (1969). An analysis of short self-report measures of life satisfaction: Correlation with rater judgments. <u>Journal of Gerontology</u>, 2, 465-469.



Yesavage, J. A., Brink, T. L., Rose, T. L., Lum, O., Huang, V., Adey, M., & Leirer, V. O. (1983). Development and validation of a geriatric depression screening scale: A preliminary report. <u>Journal of Psychiatric Research</u>, 17, 37-49.



Table 1

Correlation Matrix

	Depression	Spirituality	Hope	Hope: Agency	Hope: Pathways	Symptoms of Poor Health	Life Satisfaction
Depression	1.000			Ī			
Spirituality	558**	1.000					
Норе	600**	.488**	1.000				
Hope: Agency	657**	.466**	.891**	1.000			
Hope: Pathways	441**	.419**	.915**	.632**	1.000		
Symptoms of Poor Health	.473**	083	231	195	221	1.000	
Life Satisfaction	748**	.497**	.575**	.620**	.430**	299	1.000

^{**} Correlation is significant at the 0.01 level (two-tailed).







U.S. Department of Education

Office of Educational Research and Improvement (OERI)

National Library of Education (NLE)

Educational Resources Information Center (ERIC)

REPRODUCTION RELEASE

(Specific Document)

Title: Variables Related to Adults in Assist Author(s): Maggar-moe, JL., A			
Corporate Source: University of Kan		Publication Date:	
II. REPRODUCTION RELEASE:		Spal	ito; S.V., & Diamond, KILL
In order to disseminate as widely as possible to abstract journal of the ERIC system, <i>Resources in</i> media, and sold through the ERIC Document Regranted, one of the following notices is affixed to affirm the system of the page.	Education (RIE), are usually made availa production Service (EDRS). Credit is give each document.	able to users in microfi on to the source of eac	che, reproduced paper copy, and electronic th document, and, if reproduction release is
The sample sticker shown below will be affixed to all Level 1 documents	The sample sticker shown below will affixed to all Level 2A documents		The sample sticker shown below will be affixed to all Level 28 documents
PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL HAS BEEN GRANTED BY	PERMISSION TO REPRODUCE DISSEMINATE THIS MATERIA MICROFICHE, AND IN ELECTRONI FOR ERIC COLLECTION SUBSCRIBE HAS BEEN GRANTED BY	L IN C MEDIA	PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL IN ICROFICHE ONLY HAS BEEN GRANTED BY
TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)	TO THE EDUCATIONAL RESOU INFORMATION CENTER (ERIC)	RCES 2E	TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)
Level 1	Level 2A		Level 2B
Check here for Level 1 release, permitting reproduction and dissemination in microfiche or other ERIC archival media (e.g., electronic) and paper copy.	Check here for Level 2A release, permitting and dissemination in microfiche and in electric ERIC archival collection subscribers	onic media for	eck here for Level 2B release, permitting reproduction and dissemination in microfiche only
Doc If permission	cuments will be processed as Indicated provided repro to reproduce is granted, but no box is checked, docum	oduction quality permits. nents will be processed at Le	evel 1.
as indicated above. Reproduction from	n the ERIC microfiche or electronic media ht holder. Exception is made for non-pro	by persons other than	eproduce and disseminate these documents ERIC employees and its system contractors raries and other service agencies to satisfy
Sign here,→ Signature: Jeana J. N	Pagyon - Moe Kansas	Printed Name/Position/Title Teana L.	o: Graduate Stud Magyar-Moe M.S., Dates
please Organization/Address: University of	Kansas v #5	Telephone: (785) 2 E-Mail Address:	Magyar-Moe M.S., Doctor 86-4287 FAX: NA Date: 1/28/27



III. DOCUMENT AVAILABILITY INFORMATION (FROM NON-ERIC SOURCE):

If permission to reproduce is not granted to ERIC, or, if you wish ERIC to cite the availability of these documents from another source, please provide the following information regarding the availability of these documents. (ERIC will not announce a document unless it is publicly available, and a dependable source can be specified. Contributors should also be aware that ERIC selection criteria are significantly more stringent for documents that cannot be made available through EDRS.)

Publisher/Distributor:		
Address:		
Price:		
V. REFERRAL OF ERIC TO COPYR		
dress:		
Address:		
V. WHERE TO SEND THIS FORM:		
Send this form to the following ERIC Clearinghouse:	ERIC Counseling & Student Services University of North Carolina at Greensboro 201 Ferguson Building PO Box 26171 Greensboro, NC 27402-6171	

